

Submitted to *The Astrophysical Journal*

The Evolution of the Effective Equation of State of the IGM

Massimo Ricotti, Nickolay Y. Gnedin, and J. Michael Shull

Center for Astrophysics and Space Astronomy, Department of Astrophysical and Planetary
Sciences, University of Colorado, Campus Box 389, Boulder, CO 80309

ricotti,gnedin,mshull@casa.colorado.edu

ABSTRACT

We develop a method to extract the “effective equation of state” of the intergalactic medium from the Doppler (b) parameter distribution of the low-density Ly α forest. We test the method on numerical simulations and then apply it to published observations of the Ly α forest at redshifts $z \simeq 0$ to 4. We find that the effective equation of state is close to isothermal at redshift $z \sim 3$, indicating that a second reheating of the IGM took place at $z \sim 3$. This reheating can plausibly be identified with the reionization of He II observed to occur at $z \sim 3$.

Subject headings: cosmology: theory — intergalactic medium — equation of state — quasars: absorption lines

1. Introduction

It is widely believed that density and temperature fluctuations in the intergalactic medium (IGM) are responsible for the Ly α forest of absorption lines observed in the spectra of QSOs. This hypothesis is supported by the results of numerical simulations that are able to reproduce accurately the statistical properties and main features of the observed high-resolution spectra (cf., Weinberg *et al.* 1999 and references therein). At the same time, observations of the high redshift Ly α forest with Keck telescopes (Hu *et al.* 1995; Lu *et al.* 1996; Kirkman & Tytler 1997) and of the local Ly α forest with the Hubble Space Telescope (HST) (Shull, Penton, & Stocke 1999; Penton, Stocke, & Shull 1999) have sufficient spectral resolution and signal-to-noise to permit a detailed study of the thermal properties of the IGM and comparisons with the output of cosmological simulations.

In hierarchical dark matter dominated cosmological models, the formation of the Ly α forest depends on two loosely related pieces of information: the evolution of the dark matter and the

thermal history of the IGM.¹ Recent work by Croft *et al.* (1999) in large part solved the problem of recovering the power spectrum of the dark matter from the Ly α absorption spectra. Thus, the thermal history of the IGM remains the only piece of information still unrecovered from the observational data on the Ly α forest. This paper complements the work of Croft *et al.* (1999) by attempting to recover the thermal history of the IGM from the data.

It has been shown (Hui & Gnedin 1997) that in sufficiently low density IGM (cosmic overdensity $\delta \lesssim 5 - 10$) the gas temperature is tightly (to within a few percent) related to the gas density, with a power-law relation of the form $T = T_0(1 + \delta)^{\gamma-1}$, which we will call an “effective equation of state” because it appears to relate the density and the temperature of the gas.² The thermal history of the IGM can then be described with high precision by the time evolution of the two parameters, T_0 and γ . In the most simple physical model of the photoionized IGM adopted in all currently existing numerical simulations, this evolution is determined by the balance of adiabatic cooling due to the expansion of the universe and photoionization heating of hydrogen and helium. However, it is also possible that Compton heating from hard X-rays (Madau & Efstathiou 1999), radiative transfer effects, and time-dependent He II reionization can be important processes affecting the thermal history of the IGM (Abel & Haehnelt 1999; Ferrara & Giallongo 1999).

The most readily available piece of observational data is the joint distribution of the column densities and Doppler (b) parameters of absorption lines. Since the column density of an absorption line strongly correlates with the density of the gas in which the line originates (Miralda-Escudé *et al.* 1996; Hernquist *et al.* 1996), we can use the column density of a line as a measure of the gas density. At the same time, the width of the line contains information about the gas temperature. Thus, we can recover the information about the evolution of the “effective equation of state” of the IGM from the $b - N$ distribution of the Ly α forest. This was also noted independently by Schaye *et al.* (1999).

Undoubtedly, a better way to recover the “effective equation of state” could be developed, using the H I absorption spectra rather than $b - N$ distributions. However, this method is not yet developed. In addition, $b - N$ distributions are readily available as published data, whereas the absorption spectra are not. Thus, we proceed with analyzing $b - N$ distributions with the understanding that our results could in principle be improved upon by using the full information available in the absorption spectra.

At low redshifts ($z \sim 0$) the amount of the intergalactic gas that is shock heated is thought to be larger than at higher redshifts (Cen & Ostriker 1999). This, in principle, can induce unrecoverable systematic effects into our method. However, Davé *et al.* (1999) have shown that the “effective

¹The lack of understanding of how star formation proceeds prevents us from making a (in principle existing) direct connection between the evolution of the dark matter and the evolution of the ionizing radiation in the universe, the latter being the primary source for the evolution of the thermal state of the IGM. The QSO contribution to the ionizing radiation background that dominates up to redshift $z \sim 4$ is, instead, better known.

²The true equation of state is, of course, the ideal gas equation of state $P = nk_bT$.

equation of state” applies down to redshift $z \simeq 0$ even if the fraction of gas that is shock heated and that does not belong to the simple power-law relation increases at low redshifts. A relation between the column density and the overdensity of Ly α clouds also holds at $z = 0$. We can thus be confident that we can measure the “effective equation of state” of the IGM at low redshifts as well. Even at $z = 0$, there still exist a population of absorbers that are not shock heated and which define the lower cutoff of the $b - N$ distribution (Davé *et al.* 1999). The resulting Ly α lines are distinct, narrow features in the HST spectra (Penton *et al.* 1999); it is this population that we select by our method. In other words, the approach we adopt automatically selects against the shock-heated gas and thus is quite suitable for measuring the “effective equation of state” even at low redshift.

The paper is organized in four sections. In § 2 we describe our method to measure the “effective equation of state” of the IGM. We study the output of numerical simulations for several cosmological models in order to understand how the thermal broadening of the lines affects the $b - N$ distribution. In § 3 we describe the results of the method on the synthetic spectra created using the simulation outputs, and in § 4 we apply the method to real observations of the Ly α forest at $0 \lesssim z \lesssim 4$. In § 5 we discuss the results and their implications on understanding the thermal history of the IGM.

2. Method

We follow three logical steps to derive the “effective equation of state”. First, we analyze the result of several numerical simulations using the Voigt profile fitting code AUTOVP (Davé *et al.* 1997) to build synthetic $b - N$ distributions used to check the method. We consider several distinct cosmological models with different values of the ionizing intensity at 1 Ryd, J_{21} (in units of 10^{-21} erg cm $^{-2}$ s $^{-1}$ Hz $^{-1}$ sr $^{-1}$), and the “effective equation of state”. We also analyze different random realizations of the same model at the same or higher resolution. We use the Hydro-PM approximation to model the low-density IGM, which allows us to scan a large set of cosmological models and random realizations with sufficient numerical resolution, a task, currently unachievable with the full hydrodynamic simulations. We demonstrate that the systematic errors introduced by the HPM simulations are completely negligible compared to other systematic and statistical errors present in the method and the observational data.

We find that the thermal broadening $b_T = (13 \text{ km s}^{-1})\sqrt{T/10^4\text{K}}$ of a line with peak overdensity δ (we use the notation $\rho = \bar{\rho}(1 + \delta)$ where ρ is the baryon density and $\bar{\rho}$ is the mean baryon density) corresponds to the value of $b = b_M$ at the maximum of the $b - N$ distribution. Values of $b < b_T$ are possible because of the spread of the $N - \rho$ distribution and because the Voigt profile fitting routine introduces some lines that do not correspond to density peaks. Another reason why the cutoff of the $b - N$ distribution is not sharp is that errors on b and N smooth out the distribution. This effect is negligible for the simulated Ly α lines because we pre-determine the

continuum flux, zero flux levels and higher signal to noise, but it is important for the observational data.

Second, we find an efficient way to locate the maximum of the $b - N$ distribution. In order to use all the information available, we use the Maximum Likelihood Analysis (MLA) method to fit a parametric two-dimensional function to the data points in the $b \otimes N$ space. This allows us not only to locate the maximum of the distribution but also to study its shape. We will see that this could be a powerful way to extract information on the physics of the low-density Ly α clouds. We apply the MLA method to the simulated Ly α lines and demonstrate that the parameters T_0 and γ of the “effective equation of state” lie inside the 1σ confidence level of the values that we measure for all the models and random realizations.

Third, we apply the MLA to real data sets at redshifts $z = 4 \rightarrow 0$, and we measure the location of the maximum and the shape of the distribution. We account for the errors on b and N that, as already mentioned, spread the real shape of the distribution. At this point, we use the results of the simulations to interpret the b values at the maximum of the distribution as due to thermal broadening. This gives us the relation between T and N but not the “effective equation of state”. We still need to know the relationship between N and ρ , therefore we use the results we obtained from the simulations in conjunction with the Density Peak Ansatz approximation to determine this relationship (this is our second hypothesis based on the simulation results). It turns out that we only need to know the column density of the absorption lines with zero peak overdensity, $N_{\overline{\rho}}$, to derive the “effective equation of state” within the observational accuracy.

2.1. Simulations

We use an approximate simulation technique called HPM, developed by Gnedin & Hui (1998), to model the fluctuations in the low-density IGM ($\delta \lesssim 10$). In its essence, the HPM method uses a simple Particle-Mesh (PM) solver modified to account for the effect of gas pressure. The predefined “effective equation of state” is then used to compute the gas temperature and pressure at every point directly from the value of the cosmic gas density at this point. Thus, there is no need to introduce a special equation for the gas temperature as in a full hydrodynamic solver. As a result, the HPM approximation is only about 25% slower (due to the overhead of computing the equation of state) than a simple PM solver. It is substantially faster than a full hydrodynamic solver (due to both fewer computations at each time step and fewer time steps), while delivering results that are sufficiently accurate for our purposes.

In order to verify the accuracy of the HPM approximation, we performed Voigt profile fitting for the full hydrodynamic simulations and for the HPM simulation with the same initial conditions and resolution used in the original HPM paper (Gnedin & Hui 1998). Figure 1 shows the $b - N$ distributions for the full hydrodynamic simulation and the HPM approximation plotted on top of each other. The two distributions agree quite well.




Fig. 1.— The $b - N$ distributions for the full hydrodynamic simulation (*crosses*) and the HPM simulation (*filled triangles*) of the same Λ CDM cosmological model, the same initial conditions, and the same resolution.

Figure 2 shows the comparison between the measured b -parameters of the individual lines from two simulations. While at high b values the spread is significant, mostly due to the fact that the Doppler profile fitting is not a uniquely defined procedure, at low b values the HPM and the full hydrodynamic simulation agree on a *line-by-line basis* to better than observational uncertainties.

In order to estimate the systematic error in the position of the peak of the $b - N$ distribution introduced by the HPM simulation, we have applied our method (described below) to the two distributions shown in Fig. 1. We have found that the intercepts, b_0 , of the $b_T(N)$ relations (see equation [4]) agree within 0.5%, and the slopes, β , agree within 3.5%. Thus, the systematic error introduced by the HPM simulation is completely negligible (at least a factor of 10 smaller) compared to the other systematic and statistical uncertainties (as discussed below in more detail).




Fig. 2.— Comparison of b parameters for the corresponding lines measured in the HPM approximation versus the full hydrodynamic simulation. The low- b end of the distribution, where the maximum is located, is matched better than the observational errors.

Numerical resolution is very important for accurately modeling the Ly α forest (Gnedin 1998; Theuns *et al.* 1998; Bryan *et al.* 1999). The HPM simulations used in this paper have a cell size of $10h^{-1}$ kpc and resolve the filtering scale k_F over which the baryonic fluctuation is smoothed,

$$\frac{\delta_b}{\delta_X} \approx e^{-k^2/k_F^2},$$

(Gnedin & Hui 1998) by about a factor of three. This choice of the resolution scale has also been confirmed as sufficient by other authors using the full hydrodynamic simulations (Theuns *et al.* 1998; Bryan *et al.* 1999).

We produce synthetic spectra of the Ly α forest at $z = 2.85$ for eight of the 25 different flat cosmological scenarios simulated using the HPM approximation in Gnedin (1998). The first three models, SCDM.2A, SCDM.2D, and SCDM.2E, are three random realizations of the same model, SCDM.2L is the same cosmological model with a different “effective equation of state”, LCDM.3A

Table 1. Cosmological Models at $z = 2.85$

Model	Ω_0	Ω_b	Ω_Λ	Ω_ν	h	σ_8	n	γ	$T_{0,4}$	σ_F	J_{21}
(a) SCDM.2A	1	0.05	0	0	0.5	1	0.7	1.49	0.91	2.09	0.17
(b) SCDM.2D ^a	1	0.05	0	0	0.5	1	0.7	1.49	0.91	2.09	0.17
(c) SCDM.2E ^a	1	0.05	0	0	0.5	1	0.7	1.49	0.91	2.09	0.17
(d) SCDM.L2A ^b	1	0.05	0	0	0.5	1	0.7	1.49	0.91	2.09	0.17
(e) SCDM.2L	1	0.05	0	0	0.5	1	0.7	1.33	1.21	2.26	0.17
(f) LCDM.3A	0.35	0.03	0.65	0	0.7	1	1.04	1.51	1.02	2.27	0.30
(g) CHDM.3A	1	0.05	0	0.2	0.5	1	0.81	1.49	0.91	1.41	0.25
(h) OCDM.1A	0.3	0.04	0	0	0.7	1	0.91	1.24	2.46	1.75	0.50

^a A different random realization of the SCDM.2A model.

^b 512^3 cells and the box size of $5.12h^{-1}$ Mpc.

is a CDM model with the cosmological constant, CHDM.3A represents a class of Cold+Hot CDM models, and OCDM.1A is an open CDM model. All of these simulations have 256^3 cells and thus have a box size of $2.56h^{-1}$ Mpc. In order to test the effect of box size, we also analyze a larger simulation of the SCDM.2A model. This simulation, labeled SCDM.L2A, has 512^3 cells and box size of $5.12h^{-1}$ Mpc. As has been shown in Gnedin (1998), the $5.12h^{-1}$ Mpc size of the computational box is sufficient to reduce the cosmic variance in the column density distribution to a few percent.

The parameters of the models are summarized in Table 1, where we preserve the labeling used in Gnedin (1998). We used the AUTOVP Voigt profile fitting routine (Davé *et al.* 1997) to measure N and b for absorption lines along a thousand lines of sight in each simulation. This gives us about ten thousand absorption lines per simulation. We use this $b - N$ distribution to verify the ability of our method to recover the assumed “effective equation of state”.

2.2. Equation of State and $b - N$ Distribution

There exists a tight correlation between the intergalactic gas density and temperature in the low-density regime, where shock heating is not important (Hui & Gnedin 1997). The density-temperature relation is well described by a power-law “effective equation of state”,

$$T = T_0(1 + \delta)^{\gamma-1}, \quad (1)$$

where T_0 is the temperature of the gas at the mean cosmic density of baryons and γ describes how the temperature changes with density. It is clear that, if $\gamma = 1$, the temperature of the IGM




Fig. 3.— A scatter-plot (black dots) of the column density versus density at the peak of density fluctuations for the model LCDM.3A at $z = 2.85$, calculated by using the Density Peak Ansatz approximation and the simulations to measure the second derivative at the peak of the density perturbations (see § 3.2). The long-dashed line corresponds to the best fit using constant weighting of the lines and the solid line using weighting $\propto N^{1/2}$ to account for the 0.5 slope of the logarithmic column density distribution.

is constant everywhere (the isothermal case). The evolution of both T_0 and γ depends on the reionization history of the universe.

Thus, measuring the “effective equation of state” would potentially allow us to uncover the evolution of the ionizing background in the universe, and $b - N$ distribution of the Ly α forest offers us a way to measure T_0 and γ as a function of redshift simply because the Doppler b parameter is related (albeit indirectly due to peculiar velocities) to the gas temperature, whereas the column density of an absorber is known to be tightly correlated with the gas density.

To illustrate the latter point, we plot in Fig. 3 the column density of the absorption lines as a function of the overdensity at the peak of the line for the LCDM.3A model at $z = 2.85$ calculated by using the simulations together with the Density Peak Ansatz approximation (see § 3.2). In this




Fig. 4.— Distribution of $\log b$ versus $\log N$ for the SCDM.2D model (black dots) at $z = 2.85$. The contour plot is the $N - \rho$ distribution re-mapped on the $b \otimes N$ space using the “effective equation of state” and equation (3). This corresponds to peculiar velocities being set to zero. The solid line is the “effective equation of state” rewritten in terms of the thermal width of the lines as a function of the column density derived using equation (4) to relate the column density and the gas density. As explained in the text, we use only absorption lines in a small range of the column densities. The range of the column densities used and the number of lines are reported in Table 2.

case, a power-law,

$$N = N_{\overline{\rho}}(1 + \delta)^\Gamma, \quad (2)$$

is also a good fit to the distribution, but the correlation between N and δ is not tight and the spread of the relation depends on the cosmological model (Zhang *et al.* 1996; Hui, Gnedin, & Zhang 1997).

If we define the thermal broadening parameter,

$$b_T = \left(\frac{2kT}{m_H} \right)^{\frac{1}{2}} = (13 \text{ km s}^{-1}) T_4^{\frac{1}{2}}, \quad (3)$$




Fig. 5.— Histograms of a vertical slice at $z = 2.85$ and $\bar{N} \simeq 10^{13} \text{ cm}^{-2}$ of the scatter plot in Fig. 4. The solid line represents the b -distribution with peculiar velocities being set to zero (*i.e.*, only thermal broadening $b = b_T$) and the long-dashed line the b -distribution.

where $T_4 \equiv T/10^4\text{K}$, we can write the equation that relates the thermal width and the column density of an absorption line using (1) and (2),

$$b_T = b_0 \left(\frac{N}{N_0} \right)^\beta, \quad (4)$$

where,

$$\beta = \frac{\gamma - 1}{2\Gamma}, \quad (5)$$

$$b_0 = (13 \text{ km s}^{-1}) T_{0,4}^{1/2} \left(\frac{N_0}{N_{\bar{\rho}}} \right)^\beta, \quad (6)$$

where $T_{0,4} = T_0/10^4\text{K}$. For each simulation, the values of T_0 , γ , $N_{\bar{\rho}}$ and Γ are known (the last two are measured by fitting equation (2) to the $N - \rho$ distribution as shown in Fig. 3 and as explained in § 3.2). Therefore, using equation (4) we can derive b_T . In Fig. 4 we plot the $b - N$ distribution

for 11,520 absorption lines (black dots), taken from the spectra of the Λ CDM.2D model with $J_{21} = 0.17$ at redshift $z = 2.8$ using AUTOVP routine. The solid line is the thermal broadening b_T defined in equation (3), and the contour plot shows the $N - \rho$ distribution remapped on the $b \otimes N$ space using the “effective equation of state” and equation (3), which corresponds to peculiar velocities being set to zero. It is evident that the solid line does not define the lower envelope of the $b - N$ distribution owing to small-number statistics and to the intrinsic spread of the $N - \rho$ distribution. However, both the solid line and the lower envelope track one another. Indeed, because we decided to fit the $N - \rho$ distribution instead of finding its upper cutoff (the cutoff toward high column densities), the thermal broadening b_T will correspond to the peak of the $b - N$ distribution instead of its lower cutoff. In Fig. 5 we show this result more clearly by plotting a slice of the distribution at the mean column density $\bar{N} \simeq 10^{13} \text{ cm}^{-2}$, averaged on a bin of width $\Delta \log N \simeq 0.1$. The long-dashed line is the b -distribution and the solid line is the b -distribution with peculiar velocities being set to zero. Armed with this finding, in the next section we will seek an efficient way to locate the maximum of the $b - N$ distribution and demonstrate that, within the errors of the measurement, we are able to recover b_T .

When we apply this method to the observed Ly α forest in § 4, we can only measure directly b_T as a function of the column density N . We do not have a relation between the column density and overdensity in this case. The parameters of the “effective equation of state” are related to the measured β and $T_{0,4}^\dagger = (b_0/13 \text{ km s}^{-1})^2$ by the equations:

$$\gamma - 1 = 2\beta\Gamma, \quad (7)$$

$$T_{0,4} = T_{0,4}^\dagger \left(\frac{N_{\bar{p}}}{N_0} \right)^{2\beta}. \quad (8)$$

In § 3.2 we find an approximate relationship for Γ and $N_{\bar{p}}$ that allows us to solve the equations (7) and (8) given some cosmological parameters and the ionizing background intensity J_{21} .

2.3. Maximum Likelihood Analysis

We define a likelihood function

$$\mathcal{L} = \prod_{i=1}^n p_i, \quad (9)$$

where $p_i = p(b_i, N_i | \text{model})$ is the probability for the line i to have a width $b = b_i$ and the column density $N = N_i$ and n is the number of lines in the sample. If we describe the magnitude of the errors by a Gaussian distribution, then the probability function p_i^{obs} observed is related to the true one by the double convolution:

$$p_i^{obs} = \frac{(1 - r^2)^{-\frac{1}{2}}}{2\pi\sigma_{b,i}\sigma_{N,i}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p(b, N | \text{model}) \exp \left[- \left(\frac{\Delta b_i^2}{2\sigma_{b,i}^2} + \frac{\Delta N_i^2}{2\sigma_{N,i}^2} - r \frac{\Delta b_i \Delta N_i}{\sigma_{b,i}\sigma_{N,i}} \right) \frac{1}{1 - r^2} \right] dN db \quad (10)$$

where $\sigma_{b,i}$ and $\sigma_{N,i}$ are the errors on b_i and N_i respectively, $\Delta b_i = b_i - b$, $\Delta N_i = N_i - N$ and r is the correlation coefficient between the errors in b and N . Usually, the errors on the column density and Doppler parameter tend to be anticorrelated ($r < 0$), especially when the line is saturated, since the fit tends to preserve the equivalent width of the line. We adopt a parametric model for $p(b, N)$ and the free parameters are determined by maximizing the likelihood function (Efsthathiou *et al.* 1988).

A simple and accurate method of estimating errors is to determine numerically the ellipsoid of parameter values defined by

$$\ln \mathcal{L} = \ln \mathcal{L}_{max} - \frac{1}{2} \chi^2_{\beta}(M), \quad (11)$$

where $\chi^2_{\beta}(M) = t\sqrt{M}$ is the β -point of the χ^2 distribution with M degrees of freedom and $t\sigma$ is the confidence level wanted. The degrees of freedom are the number of lines minus the free parameters of the distribution. The parametric function, equation (13), that we adopt to fit the $b - N$ distribution, has nine free parameters, thus the ellipsoid of the errors is nine-dimensional. We are only interested in determining the errors on b_0 and β . The rigorous way to find these should be to project the hyper-ellipsoid on the $b_0 - \beta$ space but this is computationally expensive. We use an approximate method that consists of maximizing the likelihood function with respect to all the other parameters for each value of b_0 and β .

In general, the matrix of errors is not diagonal, and the errors on the parameters are correlated. The value of N_0 that makes errors on b and β uncorrelated (*i.e.*, the ellipse has principal axes parallel to the Cartesian plane and the matrix of errors is diagonal) is the average column density of the sample of lines used in the fit. We choose N_0 in order to minimize the correlation of the errors. The value of N_0 changes in each MLA mainly because the range of the column densities of the data-set changes.

These are the main advantages of the method: (i) the MLA method has well-defined asymptotic error properties (*e.g.*, Kendall & Stuart 1961); (ii) there is no requirement to select bin widths; (iii) it is possible to account for errors on both b and N ; (iv) we can also study the shape of the distribution; (v) all the data are used; (vi) there is no need to find the distribution cutoff that is not well defined because of artificial lines and metal lines. The disadvantage of this method is that it is difficult to test whether the assumed parametric form of $p(b, N)$ is a good fit to the data. In this paper, we adopt this method but we have also used a classic minimum of Chi-square algorithm to fit the function $\bar{p}(b) = p(b, N = \bar{N})$ averaged over small intervals of the column densities, selecting several bin widths of N . This preliminary study has been useful to choose the form of the parametric function that is a good fit to the data.

In the literature, the b distribution is basically fitted with two types of parametric functions: (i) a Gaussian with a low b cut-off, (*i.e.*, a three-free-parameter function) (Kirkman & Tyler 1997)

and (ii) a one-parameter function of the form (Hui & Rutledge 1999):

$$\frac{dN}{db} \propto \frac{b_\sigma^4}{b^5} \exp \left[-\frac{b_\sigma^4}{b^4} \right], \quad (12)$$

where $b_\sigma = [(b_T^2 + b_{res}^2 + b_J^2)/(D_+\sigma_0)]^{1/2}$ is the b value at the peak of the function. Here b_J is the baryon smoothing scale, b_{res} is the resolution width, D_+ is the linear growth factor ($D_+ = (z+1)^{-1}$ in a flat cosmology) and σ_0 is the rms linear density fluctuation approximately at the Jeans scale.

We find that neither of these functions provides a good fit to the data. Instead, we use equation (12) with the exponents of b in the power-law tail and in the exponential cutoff as free parameters. We notice that, as the column density increases, the power-law tail of the distribution becomes more narrow and the exponential cutoff becomes more sharp. We have also tried a simple power-law b^{-n} with a low b cutoff, motivated by the fact that the exponential cutoff is very sharp, but we have found that this form of the distribution is not a good fit to the data.

The form of the parametric function that is the best fit to the simulated Ly α forest and that we will use in the rest of the paper is,

$$p(b, N) = \mathcal{N} N^\alpha \left(\frac{b_*}{b} \right)^\varphi \exp \left[-\left(\frac{b_*}{b} \right)^\Phi \right], \quad (13)$$

where,

$$\begin{aligned} \varphi &= \varphi_0 + \varphi_1 \left(\frac{N}{N_0} \right) + \varphi_2 \ln \left(\frac{N}{N_0} \right), \\ \Phi &= \Phi_0 + \Phi_1 \left(\frac{N}{N_0} \right) + \Phi_2 \ln \left(\frac{N}{N_0} \right), \\ b_* &= b_M \left(\frac{\varphi}{\Phi} \right)^{\frac{1}{\Phi}}, \end{aligned} \quad (14)$$

$$\mathcal{N} = \left(\frac{\alpha - 1}{N_{min}^{1-\alpha} - N_{max}^{1-\alpha}} \right) \left(\frac{\Phi}{b_* \Gamma \left[\frac{\varphi-1}{\Phi} \right]} \right). \quad (15)$$

Here, $\mathcal{N}$ is the normalization constant, N_{min} and N_{max} are the low and high column density cutoffs of the distribution and $\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt$ is the Gamma function. The b value at the maximum of the function is b_M . As previously stated, we assume $b_T = b_M$. We do not use the Hui & Rutledge relation for b_σ because the Jeans smoothing scale in this expression is needed only if we want to relate the dark matter density to the baryon density and temperature. We already include this smoothing scale in the $N(\rho)$ expression that we derive in § 3.2. We assume that the broadening of the lines due to the simulation/observation resolution is negligible in the range of the column densities considered. We also find that the values of the parameters Φ_1 , Φ_2 and φ_2 are practically zero in all the MLAs for the simulations and observations as well. The shape of the distribution is then determined by 6 parameters: b_0 and β which determine the value of b_M , φ_0




Fig. 6.— Value of the free parameters in equation (13) as a function of the correlation coefficient r . The parameters are normalized to their value at $r = 0$.

and φ_1 which are related to the power law tail, Φ_0 related to the exponential cutoff, and α the slope of the column density distribution.

We do not know the correlation coefficient r of the errors on b_i and N_i . However, as is shown in Fig. 6, when we change r between zero and -0.95, the values of the parameters determined with the MLA change by at most 20%, well below the observational accuracy or the accuracy of our method.

3. Testing the Method

Table 2. Simulated Ly α forest

Model	number of lines ^c	$\log N_{min}$ (cm ⁻²)	$\log N_{max}$ (cm ⁻²)	$\log N_0$ (cm ⁻²)	b_0 (km s ⁻¹)	β	φ_0	φ_1	Φ_0	α
(a) SCDM.2A	11500/1716	13.2	14.6	13.7	16.8	0.18	3.1	3.5	3.6	1.9
(b) SCDM.2D ^a	11520/1744	13.2	14.6	13.7	18.0	0.16	3.9	3.7	3.8	1.8
(c) SCDM.2E ^a	10222/1204	13.2	14.6	13.7	18.5	0.15	4.0	3.7	3.7	1.9
(d) SCDML.2A ^{a,b}	12131/1373	13.3	14.5	13.7	19.0	0.16	4.1	3.1	2.5	1.8
(e) SCDM.2L	6980/1370	13.0	14.9	13.7	19.2	0.11	6.9	3.7	2.2	1.7
(f) LCDM.3A	9233/1182	13.2	14.4	13.7	19.4	0.14	3.6	3.7	4.9	1.7
(g) CHDM.3A	11812/1496	13.2	14.5	13.7	17.6	0.17	2.8	3.7	4.2	1.9
(h) OCDM.1A	4289/271	13.5	14.9	13.7	26.1	0.04	12	0.1	13	1.9

^a A different random realization of the SCDM.2A model.

^b 512^3 cells and the box size of $5.12h^{-1}$ Mpc.

^c Total number of lines/number of lines in the column density range $N_{min} \leq N \leq N_{max}$.

3.1. The $b - N$ Distribution

The number of absorption lines in each simulation is about 10^4 , but we are restricted to use lines in a range of the column densities approximately one decade wide. This reduces the number of lines to about 10^3 . The upper limit on N is determined for each simulation by imposing a maximum overdensity at the line center, $\delta \sim 10$, which is the limit below which HPM simulations produce accurate results. The lower limit on the column density is determined by the completeness requirement of the distribution and by intrinsic problems associated with the Voigt fitting routine AUTOVP. The relative errors on b and N are about 10%, smaller than typical observational errors, since we assumed a higher signal-to-noise ratio than is usually afforded by the observational data (in order to test our method in more challenging conditions) and because we do not have uncertainties due to continuum fitting and zero flux level. Therefore, the convolution with Gaussian errors does not affect the shape of the distribution in a significant way.

In Table 2 we show the values of the parameters that give the best fit to the $b - N$ distribution. We note that the value of φ_1 is surprisingly constant for all the models but one. Also, the lack of lines in the region of large N and b turns out to be a real feature of the distribution and not an apparent feature due to a decreasing number of lines. The width of the b distribution at fixed N is proportional to $1/\varphi$ and decreases with N as $1/(\varphi_0 + \varphi_1 N)$. The value of Φ , instead, varies between four and six and is directly related to the width of the $N - \rho$ distribution.

In Fig. 7 we present the result with which we are more concerned in this paper: that the parameters of the equation of state can be measured from the $b - N$ distribution within the




Fig. 7.— Error ellipses for the models at $z = 2.85$, listed in Table 1. The solid lines show 1σ error contours and the dashed lines show the 2σ error contours. The crosses are the expected values of b_0 and β calculated with equation (3). The size of the crosses corresponds to the systematic error introduced by the HPM approximation.

precision of the method. Each panel shows the best fit (black dot) and the 1σ and 2σ error contours for b_0 and β . The triangle and the cross are the theoretical values derived using equation (4) for two slightly different measurements of the parameters $N_{\bar{\rho}}$ and Γ in the equation (2). In one case, we fit the $N - \rho$ distribution with constant weighting for all the lines (triangles), and in the other (crosses) with a $N^{1/2}$ weighting in order to correct for the effect of the decreasing number of lines with increasing the column density. For a more detailed explanation see § 3.2. In all the models, the expected values of b_0 and β lie near the 1σ contour level of the measurement, as they should be if the errors are estimated correctly.

3.2. Relation Between Column Density and Overdensity

The Density Peak Ansatz (DPA) approximation is based on the assumption that a single Ly α line arises from a peak in the cosmic gas density along the line of sight to a distant quasar. Then, given the value of the cosmic density at the peak, and the second derivative of the density along the line of sight (the first derivative vanishes at the peak), a value for the column density can be associated with the peak as follows (Gnedin & Hui 1996):

$$N_{\text{HI}} \equiv (3.35 \times 10^{13} \text{ cm}^{-2}) T_{0,4}^{-0.7} \left(\frac{\Omega_b h^2}{0.019} \right)^2 \left(\frac{0.5}{J_{21}} \right) \left(\frac{1+z}{4} \right)^5 \times (1+\delta)^{2-0.7(\gamma-1)} \left[-(1-0.35(\gamma-1)) \frac{d^2 \ln(1+\delta)}{dx^2} \right]^{-1/2}, \quad (16)$$

and x being measured in Mpc. Here J_{21} , is the ionizing intensity,

$$J_{21} \equiv \frac{\int J_\nu \sigma_\nu d\nu / \nu}{\int \sigma_\nu d\nu / \nu} \times \frac{1}{10^{-21} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}},$$

where J_ν is the radiation specific intensity and σ_ν is the HI photoionization cross section. Comparing equations (2) and (16) we find that one can write the inverse of the square root of the average curvature at a density peak, *i.e.* the typical comoving size (expressed in Mpc) of a Ly α cloud, as a function of the overdensity,

$$\left(-\frac{d^2 \ln(1+\delta)}{dx^2} \right)^{-1/2} = \Xi (1+\delta)^\xi, \quad (17)$$

where Ξ and ξ are model dependent. We hypothesize that the average curvature of a density peak is a function of the non-linearity of the baryonic density field. An accurate indicator of the non-linearity of the gas distribution of a given cosmological model is the linear density fluctuation at the filtering scale, σ_F (Gnedin & Hui 1998). We therefore fitted Ξ and ξ as linear function of σ_F for 25 flat cosmological models (Gnedin 1998). The result of the fit is given in Table 3. The parameters in equation (2) are measured using the minimum of Chi-square analysis of the $N - \rho$ distribution with a weighting on each line $\propto N^{1/2}$ in order to account for the 0.5 slope of the logarithm of the column density distribution.

Combining equations (17) and (16), we write the relationships,

$$\Gamma = 2 - 0.7(\gamma - 1) + \xi \simeq 1.68 + 0.068\sigma_F - 0.7(\gamma - 1),$$

$$N_{\overline{\rho}} = (3.35 \times 10^{13} \text{ cm}^{-2}) T_{0,4}^{-0.7} \left(\frac{\Omega_b h^2}{0.019} \right)^2 \left(\frac{0.5}{J_{21}} \right) \left(\frac{1+z}{4} \right)^5 \frac{0.19 - 0.025\sigma_F}{[1 - 0.35(\gamma - 1)]^{1/2}}. \quad (18)$$

Both Γ and $N_{\overline{\rho}}$ are insensitive to the value of σ_F if we make a conservative assumption that it is in the range 2 ± 1 (Gnedin 1998). From now on, we use a value $\sigma_F = 2$.

Table 3. Parameters of the best fit

	a	b	χ^2
$\Xi = a + b\sigma_F$	0.19 ± 0.02	-0.025 ± 0.01	0.005
$\xi = a + b\sigma_F$	-0.319 ± 0.02	0.068 ± 0.01	0.013

Finally, if we define $N_{\bar{\rho}}^* = N_{\bar{\rho}} T_4^{0.7}$, we can write the equations that relate the measured parameters β and $T_{0,4}^\dagger$ to the “effective equation of state” parameters,

$$\gamma - 1 \simeq \frac{3.634\beta}{1 + 1.4\beta}, \quad (19)$$

$$T_{0,4} = \left[T_{0,4}^\dagger \left(\frac{N_{\bar{\rho}}^*}{N_0} \right)^{2\beta} \right]^{\frac{1}{1+1.4\beta}}, \quad (20)$$

where,

$$\frac{N_{\bar{\rho}}^*}{N_0} \simeq \frac{(4.7 \times 10^{12} \text{ cm}^{-2})}{N_0} \left(\frac{\Omega_b h^2}{0.019} \right)^2 \left(\frac{0.5}{J_{21}} \right) \left(\frac{1+z}{4} \right)^5 \sqrt{\frac{1 + 1.4\beta}{1 + 0.128\beta}}. \quad (21)$$

Propagating the errors on β and b_0 , we write the errors on the “effective equation of state” parameters,

$$\frac{\Delta(\gamma - 1)}{\gamma - 1} = \frac{1}{1 + 1.4\beta} \frac{\Delta\beta}{\beta}, \quad (22)$$

$$\frac{\Delta T_{0,4}}{T_{0,4}} \simeq \frac{2}{1 + 1.4\beta} \left(\left| \frac{\Delta b_0}{b_0} \right|^2 + \left| \left(\ln \frac{N_{\bar{\rho}}^*}{N_0} - 1.4 \ln \frac{b_0}{13} \right) \frac{\Delta\beta}{1 + 1.4\beta} \right|^2 \right)^{\frac{1}{2}}. \quad (23)$$

If $N_{\bar{\rho}}^*/N_0 = (b_0/13 \text{ kms}^{-1})^{1.4}$, the errors are uncorrelated and $\Delta T_{0,4}$ has a minimum value.

4. Observed Evolution of the Equation of State

4.1. Observations

We apply our method to the published data sets of Ly α absorption lines. In this paper, we analyze eight data sets, but the “effective equation of state” can be measured with much better accuracy if we use more observations. In order to obtain good results, we need to use the data sets with small errors on b and N and with as many lines as possible inside the range of the column densities where the completeness of the sample is good. For the high-redshift Ly α forest

Table 4. Observational data

QSO	Telescope	FWHM km s ⁻¹	S/N	Voigt fitting code	number of lines
Q0000-26	Keck HIRES	6.6	20-80	VPFIT	337
APM 08279+5255	Keck HIRES	6	80	...	380
Hu <i>et al.</i> ^a	Keck HIRES	8	50	...	1056
GB 1759+7539	Keck HIRES	7	25	...	318
HS 1946+7658	Keck HIRES	7.9	15-100	VPFIT	459
B2 1225+317	Kitt Peak 4-m	18	2-20	...	159
Penton <i>et al.</i> ^b	HST/GHRS	19	10-42	...	80

Note. — The data for the quasars at high- z are from Lu *et al.* 1996 (Q0000-26), Ellison *et al.* 1999 (APM 08279+5255), Outram *et al.* 1999 (GB1759+7539), Kirkman & Tytler 1997 (HS 1946+7658), Khare *et al.* 1997 (B2 1225+317).

^aCollection of high- z quasars Q0014+813, Q0302-003, Q0636+680, Q0956+122 (Hu *et al.*, 1995).

^bCollection of low- z quasars: AKN120, ESO141-G55, FAIRALL9, MARK279, MARK290, MARK421, MARK509, MARK817, MARK501, PKS2155, Q1230+0115, H1821+643, ESO141-G55 (Penton *et al.* 1999).

we use Keck-HIRES observations of the quasars Q0000-26 (Lu *et al.* 1996) and APM 08279+5255³ (Ellison *et al.* 1999) at $z \sim 4$, Q0014+813, Q0302-003, Q0636+680, Q0956+122 (Hu *et al.* 1995), GB1759+7539 (Outram *et al.* 1999) and HS 1946+7658 (Kirkman & Tytler 1997) at $z \sim 3$. At intermediate redshift we use the data-set of Khare *et al.* 1997 for the quasar B2 1225+317 at $z \sim 1.9$. Finally, we use HST observations of the local Ly α forest kindly provided by Penton, Stocke, & Shull (1999), using the Goddard High Resolution Spectrometer (GHRS) with 19 km s⁻¹ resolution.

We use a subsample of 337 Ly α absorption lines of the complete Lu *et al.* (1996) data set within a redshift range $3.425 \leq z \leq 3.977$. The upper cutoff is chosen because this is where the UV ionizing intensity from the quasar is expected to roughly equal that of the UV background.

³The data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation.




Fig. 8a




Fig. 8b

Fig. 8.— (a) Distribution of $\log b$ versus $\log N$ for $\text{Ly}\alpha$ lines for Q0000-26 at $\langle z \rangle \sim 3.7$, with errorbars from VPFIT. (b) Comparison between the b distribution for Q0000-26 (long-dashed line) and for model LCDM.3A (solid line) at $z = 2.8$ and $\bar{N} \simeq 10^{13} \text{ cm}^{-2}$. The large errors on b and N of the observed $\text{Ly}\alpha$ forest spread considerably the true distribution.

The data set from Hu *et al.* (1995) is a collection of four quasars and consists of around 10^3 lines in the redshift range $3.425 \leq z \leq 3.977$. Unfortunately, the published list of lines does not have errors on the b parameter and the column density. In order to use this large list of lines, we decided to produce synthetic errors, assuming that their amplitude and distribution follows the same statistics as the Lu *et al.* (1996) sample. We chose this sample because it has similar resolution and S/N. Unfortunately, the Voigt profile fitting routine is not the same.

The local $\text{Ly}\alpha$ data set (Penton *et al.* 1999) consists of 79 lines from 15 Seyfert galaxies and quasars detected in observations with the HST/GHRS, using both pre- and post-COSTAR optics. We do not use the pre-COSTAR observations because this sample appears to have systematically larger b -values due to spacecraft wobble (Penton *et al.* 1999). This reduces the sample to 43 lines, which are distinct $\text{Ly}\alpha$ absorption features at $> 4\sigma$ significance, with $\langle b \rangle = 35.0 \pm 16.6 \text{ km s}^{-1}$. At low redshift, a typical line of sight to a QSO at $z \sim 0.05$ may have fewer than 10 absorbers. As a result, it is not possible to have an uniform sample of lines belonging to the same line of sight such as for the high-redshift $\text{Ly}\alpha$ forest. However, the ensemble of low- z $\text{Ly}\alpha$ absorbers, drawn from 15 sight-lines, adequately characterizes the distribution of b -values down to the 19 km s^{-1} spectral resolution of the HST/GHRS. Additional $\text{Ly}\alpha$ lines are currently being observed with the Space Telescope Imaging Spectrograph (STIS) and will double the sample of lines.

In all the samples, we do not include identified metal lines. We only use absorption lines in a limited range of the column densities. The lower limit is determined by imposing the following two conditions:

- the completeness of the sample must be close to one;
- the width of the b distribution must be monotonically decreasing in the range considered.

The first condition is usually the most stringent. In the case of Q0000-26, the two conditions are comparable. The upper limit on the column density is set by imposing the following constraints:

- overdensity of the line $\delta \lesssim 10$ (range of validity of the HPM approximation).
- $N_{max} \leq 10^{14.5} \text{ cm}^{-2}$ in order to avoid heavily saturated lines.

By restricting ourselves to this range of the column densities, we are more confident of the reliability of the results from the Voigt profile fitting algorithm. Indeed, we find that the shapes of the b -distributions obtained by different authors, using different Voigt profile fitting algorithms, are very similar in this range of the column densities. It is however possible that fluctuations of the ionizing background (Haardt & Madau 1996; Fardal *et al.* 1998) can produce different mean absorption of the quasar spectra and result in a spatially fluctuating value of $N_{\overline{\nu}}$. However, this effect is much smaller than the errors of the measurement. We will quantify the change of the IGM temperature produced by a change in the ionizing background below.

We discard lines with errors on b or N greater than 50% in order to obtain more precise results. Large errors degrade the precision of the final result. On the other hand, the number of lines of the sample decreases if we consider only the lines with small errors. By performing several MLAs, with various error thresholds, we find that limiting the relative errors to about 50% is the best choice.

In Table 4 we summarize the main properties of the data sets used in this paper. To give an idea of the magnitude of the errors and how they can affect the shape of the b distribution, we show in Figs. 8a and 8b the data set of Lu *et al.* (1996), with error-bars and the b distribution at $\overline{N} \sim 10^{13} \text{ cm}^{-2}$ compared to a typical simulated b distribution.

4.2. Results

In Table 5 we report the parameters for the best fit from the MLA. We use the same form of the function as adopted in the simulations (see equation [13]). Fig. 9 shows the values of b_0 and β and the error contours at 1σ and 2σ confidence levels. Each panel shows the results for a different quasar or data set. In Table 5 the quasar and the mean redshift of the Ly α forest are reported along with the label that appears on the upper right corner of each panel. Fig. 10 is analogous to

Table 5. Observed Ly- α forest

QSO	$\langle z \rangle$	number of lines ^c	$\log N_{min}$ (cm ⁻²)	$\log N_0$ (cm ⁻²)	b_0 (km s ⁻¹)	β	φ_0	φ_1	Φ	α
(a) Q0000-26	3.7	195	13.5	14.15	23.6	0.11	3.6	0.1	4.6	1.7
(b) APM 08279+5255	3.42	186	13.0	13.7	23.1	0.09	2.6	0.3	5.6	1.7
(c) Hu <i>et al.</i> ^a	2.85	636	13.0	13.7	24.3	0.04	4.6	0.01	3.1	1.6
(d) Q0014+813	2.95	159	13.0	13.7	26.4	0.07	5.9	0.1	2.1	1.7
(e) Q0302-003	2.85	160	13.0	13.7	26.9	0.11	4.8	0.2	2.8	1.7
(f) Q0636+680	2.75	193	12.8	13.7	23.9	0.05	4.7	0.06	3.3	1.6
(g) Q0956+122	2.85	180	12.8	13.7	25.7	0.04	4.3	0.02	4.4	1.6
(h) GB 1759+7539	2.72	13.5	13.9	13.7	23.2	0.16	3.1	1.5	2.0	2.1
(i) HS 1946+7658	2.7	221	13.0	13.6	22.4	0.06	3.3	0.05	7.2	1.8
(l) B2 1225+317	1.9	99	13.2	13.7	24.4	0.1	6.2	0.5	1.2	1.8
(m) Penton <i>et al.</i> ^b	0.06	43	12.5	13.48	34.2	0.35	12.4	0.7	1.4	1.6

^{a,b} See notes on Table 4.

^c Number of lines used in the MLA in the column density range $N_{min} \leq N \leq N_{max}$.

Fig. 9 but shows the value and error contours for $\gamma - 1$ and $T_{0,4}^\dagger$ derived using equation (19). Finally in Fig. 11 we show the values and the 1σ error contour for $\gamma - 1$ and T_0 (computed with equation [20]). In each panel we have the result for two different values of the ionizing intensity J_{21} . The values chosen for J_{21} at different redshifts are based on observational (Lu *et al.* 1996; Shull *et al.* 1999) and theoretical (Haardt & Madau 1996; Fardal *et al.* 1998; Valageas & Silk 1999) results. We adopt as minimum values of J_{21} : $J_{21}^{min}(z = 4) = 0.2$, $J_{21}^{min}(z = 3) = 0.5$, $J_{21}^{min}(z = 2) = 0.2$ and $J_{21}^{min}(z = 0) = 0.01$. The maximum values are, $J_{21}^{max}(z) = 2 \times J_{21}^{min}(z)$ for all z .

The error ellipses are strongly elongated and tilted with respect to the Cartesian axis. This happens because $N_\rho^* \neq N_0$. In this case, as shown in equation (23), the error on T_0 depends on the value of $\gamma - 1$. It is impossible to remove this correlation of the errors by changing the value of the normalization parameter N_0 because this would introduce exactly the same amount of correlation from the errors on b_0 and β due to the limited range of the column densities of the lines. Using a smaller value of the minimum column density of the distribution of absorption lines would be the only way to reduce this correlation. This means that, in order to have $N_\rho^*/N_0 = 1$, we would need a data set that is also complete for lines arising from the density fluctuations in the underdense regions ($\delta < 0$). This would probably imply the use of a more complicated form of the parametric function if the b distribution starts to become narrower in underdense regions as can be expected on theoretical grounds.

In Fig. 12, we show the final result of this work, the evolution of the “effective equation of state” of the IGM with 1σ error-bars and for two different values of the ionizing radiation J_{21} ,




Fig. 9.— Error ellipses in the $\beta \otimes b_0$ space for the data sets listed in Table 5. Solid lines show 1σ error contours and the dashed lines show the 2σ error contours.

chosen as explained above. The results are also listed in Table 6 and Table 7 for reference. The point at $z \sim 4$ is the weighted average of the observations (a) and (b), and the point at $z \sim 3$ is the weighted average of the observations (d), (e), (f), (g), (h), (i), where the letters refer to the list in Table 5.

The lines in Fig. 12 show simple photoionization models for the thermal history of the universe computed using the method of Hui & Gnedin (1998). In all the models, hydrogen is reionized suddenly at $z = 7$. The solid line shows the model where He II is not reionized at all, the short-dashed lines show models with sudden helium reionization at $z = 3$, the long-dashed line shows the same model but with the He II photoheating rate artificially increased by a factor of four following the suggestion by Abel & Haehnelt (1999). Finally, the dot-dashed line shows a model where the He II reionization is more gradual, and it provides a remarkably good fit to the data.




Fig. 10.— Error ellipses in the $\gamma - 1 \otimes \log T_0^\dagger$ space for the data sets listed in Table 5. Solid lines show 1σ error contours and the dashed lines show the 2σ error contours.

Table 6. Time evolution of the observed “effective equation of state”.

$\langle z \rangle$	$\gamma - 1$	$T_0/10^4 \text{ K}$	$T_0/10^4 \text{ K } (J_{21} \times 2)$	$\Delta T_0/T_0$	J_{21}
3.56 ± 0.27	0.38 ± 0.18	1.98	1.72	16%	0.2
2.75 ± 0.27	0.22 ± 0.10	2.52	2.21	9%	0.5
1.90 ± 0.30	0.32 ± 0.30	1.77	1.57	32%	0.2
0.06 ± 0.01	0.85 ± 0.30	0.47	0.34	37%	0.01




Fig. 11.— 1σ error ellipses in the $\gamma - 1 \otimes \log T_0$ space for the data sets listed in Table 5. The two ellipses in each panel are calculated using $J_{21} = J_{21}^{min}, J_{21}^{max}$. Values are reported in the text.

4.3. Sanity Check

We would like to emphasize that the thermal history of the universe which we derive from the observational data is quite different from the one assumed in all the simulations we used to test the method: the temperature is significantly higher and the “effective equation of state” flattens out at $z = 3$. As a sanity check, and in order to test the reliability of the method for the thermal history similar to the one observed, we ran a simulation of a “realistic” LCDM model with $\Omega_0 = 0.3$, $\Omega_\Lambda = 0.7$, $\Omega_b = 0.04$, $h = 0.7$, $n = 1$, $\sigma_8 = 0.91$ and an equation of state shown by the long-dashed line in Fig. 12, which is sufficiently similar to the observational data. We follow exactly the same procedure used to analyze the observations to derive T_0 and γ .

Fig. 13 demonstrates that our method works in this case as well, despite the fact that it was developed, based on the simulations with the thermal history different from the one observed. In that figure, we show the “realistic model” over mentioned at redshifts $z = 3.69, 3.34, 2.85, 2.31$, as




Fig. 12a




Fig. 12b

Fig. 12.— The evolution of γ (a) and T_0 (b) with redshift. The point with error bars at $z \sim 4$ is the weighted average of the observations (a) and (b), and the point at $z \sim 3$ is the weighted average of the observations (d), (e), (f), (g), (h), (i), where the letters refer to the list in Table 5. The solid line represents the theoretical model with a sudden H I reionization at $z_{\text{rei}} = 7$ and with $T_{\text{rei}} = 2.5 \times 10^4$ K. The dashed lines show the effect of sudden He II reionization in the optically thin approximation (*short-dashed line*) and with the He II photoheating rate quadrupled (*long-dashed line*). The model shown by the dot-dashed line has a quadrupled He II heating rate and a more gradual He II reionization.

well as the assumed thermal history (dashed line). One can see that our method reproduces the correct answer within the errors.

Comparing Table 2 with Table 5, we note that the simulated and observed $b - N$ distributions are very similar apart from the values of the parameter φ_1 . In all simulations but one, the parameter φ_1 , which measures how much the b -distribution narrows at high column densities, is about 3.7. In the observational data, it is indistinguishable from zero, which implies that the observed b -distribution has the same width for all values of the column density. In order to understand this difference between the b -distributions we applied our method to a sample of 60 $\text{Ly}\alpha$ lines at $z = 3$ from a cosmological hydrodynamic simulation of Miralda-Escudé *et al.* (1996), where gas shock heating is included. For this simulation we found a value $\varphi_1 = 0.1$, consistent with the observations. Our interpretation of this result is that shock heating spreads the b -distribution at high column densities. This does not affect in any way the validity of our determination of the

Table 7. Measured “effective equation of state” of the observations at $z \sim 4$ and $z \sim 3$.

$\langle z \rangle$	$\gamma - 1$	$T_0/10^4$ K	$T_0/10^4$ K ($J_{21} \times 2$)	$\Delta T_0/T_0$	J_{21}
3.70 ± 0.27 (a)	0.35 ± 0.2	2.04	1.78	15%	0.2
3.42 ± 0.27 (b)	0.47 ± 0.4	1.78	1.51	25%	0.2
2.95 ± 0.25 (d)	0.23 ± 0.3	2.69	2.47	30%	0.5
2.85 ± 0.25 (e)	0.34 ± 0.3	2.21	1.94	30%	0.5
2.75 ± 0.25 (f)	0.17 ± 0.2	2.43	2.27	20%	0.5
2.85 ± 0.25 (g)	0.14 ± 0.3	2.97	2.82	25%	0.5
2.72 ± 0.33 (h)	0.30 ± 0.3	1.77	1.58	35%	0.5
2.70 ± 0.30 (i)	0.14 ± 0.25	2.69	2.56	20%	0.5




Fig. 13a




Fig. 13b

Fig. 13.— The evolution γ (a) and T_0 (b) as a function of redshift for the “realistic” LCDM model with sudden He II reionization at $z \sim 3$ and a photoheating rate 4 times the optically thin approximation (*dashed line*).

“effective equation of state”, but instead can, in principle, be used to measure the fraction of the gas at a given density that is shock-heated.

5. Summary and Discussion

We note two main features of Fig. 12: the value of $\gamma - 1$ at $z \sim 3$ is very low and the temperature at $z \sim 3$ is too high. These features are only marginally consistent with simple photoionization models in which H I and He II are photoionized simultaneously at $z \sim 5$. The direct interpretation of this disagreement is that H I reionization occurs quite late, after $z = 5$ or so. This, however, contradicts the observation of Ly α emission lines from quasars and galaxies at $z > 5$ (Weymann *et al.* 1999; Chen *et al.* 1999), which would be impossible if the universe was neutral at $z \gtrsim 5$ (Miralda-Escudé & Rees 1998). Another interpretation of our results is that the universe reheated a second time at $z \sim 3$. Reheating will increase the temperature of the IGM and flatten the effective equation of state ($\gamma - 1$ decreases). One of the most plausible source of the secondary reheating is He II reionization. Recent observations of the He II Ly α forest also support the conclusion that He II reionization occurs at $z \sim 3$ (Reimers *et al.* 1997; Anderson *et al.* 1999; Heap *et al.* 1999). Abel & Haehnelt (1999) have recently shown that radiative transfer effects can increase the He II photoheating rate respect to the optically thin approximation by up to a factor 4. This results in an increase of the temperature by a factor 1.5-2.5.

The dashed lines in Fig. 12 show the effect of a sudden He II reionization at $z = 3$ in the optically thin approximation (the short-dashed line) and with the He II photoheating rate quadrupled (the long-dashed line). The model shown by the dot-dashed line has the quadrupled He II heating rate and a somewhat gradual He II reionization. Despite being so simple-minded, the latter model provides a remarkably good fit to the observed thermal history of the IGM.

Other possible sources of heating in the IGM are photoelectric dust heating (Nath, Sethi, & Shchekinov 1999) and Compton heating by hard X-rays. Recent results (Madau & Efstathiou 1999) indicate that the second could be an important heating source at redshifts larger than $z \sim 10$ but photoheating is dominant at lower redshifts. Photoelectric heating of dust grains could be a minor source of heat but some recent observations show only a small amount of dust in the high- z Ly α forest (Outram *et al.* 1999).

We would like to emphasize here that, while our error bars on the thermal history of the IGM are quite large, a substantial improvement can be obtained with a larger data set and extending the completeness of the sample to low- N lines. The error bars are still dominated by the statistical errors, as can be seen from comparing Fig. 7 and Fig. 9. With a larger data set, the errors on T_0 and γ can be reduced up to a factor of three, after which the systematic errors become significant. The most significant systematic error is the lack of knowledge of the intrinsic shape of the $b - N$ distribution; with the larger data set this error can also be reduced by introducing a more complicated fit to the overall distribution. The errors on the parameters β and b_0 measured with

the MLA method are proportional to the square root of the number of lines of the sample. For a typical simulation with 10^3 lines, we have $\Delta\beta \sim 0.06$ and $\Delta b_0 \sim 1.5 \text{ km s}^{-1}$. For the Hu *et al.* (1995) with 636 lines, we have $\beta \sim 0.08$ and $\Delta b_0 \sim 2 \text{ km s}^{-1}$. The errors on the “effective equation of state” parameters are approximately $\Delta(\gamma - 1) \lesssim 3.634\Delta\beta$ and $\Delta T_0/T_0 \gtrsim 2\Delta b_0/b_0$. At redshift $z \sim 3$ we have six QSOs spectra and the errors are $\Delta(\gamma - 1) \sim 0.1$ and $\Delta T_0/T_0 \sim 9\%$. Thus, more observations at redshift $z \sim 4$ and $z \sim 2$ are needed to reduce the current errors. Reducing the errors on the Doppler parameters and the column densities of the lines are important as well, in order to obtain an accurate measure of the “effective equation of state” and of the shape of the $b - N$ distribution. Also, more absorption lines for the local Ly α forest are required in order to put strong constraints on the evolution of the “effective equation of state” of the IGM at low z .

One of the major issues for future work concerns the distribution of Doppler b -values for the low- z Ly α absorbers (Penton et al. 1999). Although we worked here with just 43 absorption lines at $z < 0.06$, many more lines of sight are currently being analyzed with HST/STIS spectroscopic data. Thus, theoretical predictions of the thermal evolution of the IGM at low redshift will soon be testable. One of the intriguing predictions of many hydrodynamic simulations (cf., Cen & Ostriker 1999) is that the relative fractions of warm (photoionized) absorbers and hot (shocked) shift considerably between $z = 2$ and $z = 0$. The empirical distribution of b parameters for Ly α absorbers (from HST) and Ly β absorbers (from FUSE) should reflect this shift and can provide a test of these models.

We are grateful to Steve Penton for the as yet unpublished data set on the local Ly α forest and to Mark Fardal for the list of published data on the high-redshift Ly α forest. We thank David Tytler, Len Cowie, and Limin Lu for allowing us to use the published $b - N$ distributions. We also thank Romeel Davé for letting us use his AUTOVP automated Voigt profile fitting software. This paper was significantly improved as the result of fruitful conversations with Andrew Hamilton and Jordi Miralda-Escudé. This work was supported in part by grants from NASA (NAG5-7262) and NSF (AST96-17073). Calculations were performed the NCSA Origin2000 array under the grant AST-960015N.

REFERENCES

- Abel, T., & Haehnelt, M. G. 1999, ApJ, 520, L13
- Anderson, S. F., Hogan, C. J., Williams, B. F., & Carswell, R. F. 1998, AJ, 117, 56
- Bryan, G. L., & Machacek, M. 1999, ApJ submitted (astro-ph/9906459)
- Bryan, G. L., Machacek, M., Anninos, P., & Norman, M. L. 1999, ApJ, 517, 13
- Cen, R., & Ostriker, J. P. 1999, ApJ, 519, L109
- Chen, H.-W., Lanzetta, K. M., & Pascarelle, S. 1999, Nature, 398, 586

- Croft, R. A. C., Weinberg, D. H., Pettini, M., Hernquist, L., & Katz, N. 1999, *ApJ*, 520, 1
- Davé, R., Hernquist, L., Weinberg, D. H., & Katz, N. 1997, *ApJ*, 477, 21
- Davé, R., Hernquist, L., Katz, N., & Weinberg, D. H. 1999, *ApJ*, 511, 521
- Efstathiou, G., Ellis, R. S., & Peterson, B. A. 1988, *MNRAS*, 232, 431
- Ellison, S. L., Lewis, G. F., Pettini, M., Sargent, W. L. W., Chaffee, F. H., Foltz, C. B., Rauch, M., & Irwin, M. J. 1999, *PASP*, 111, 946
- Fardal, M. A., Giroux, M. L., & Shull, J. M. 1998, *AJ*, 115, 2206
- Ferrara, A., & Giallongo, E. 1996, *MNRAS*, 282, 1165
- Gnedin, N. Y. 1998, *MNRAS*, 299, 392
- Gnedin, N. Y., & Hui, L. 1996, *ApJ*, 472, L73
- Gnedin, N. Y., & Hui, L. 1998, *MNRAS*, 296, 44
- Haardt, F., & Madau, P. 1996, *ApJ*, 461, 20
- Heap, S. R., Williger, G. M., Smette, A., Hubeny, I., Sahu, M., Jenkins, E. B., Tripp, T. M., & Wikler, J. N. 1999, *ApJ*, submitted (astro-ph/9812429)
- Hernquist, L., Katz, N., Weinberg, D. H., & Miralda-Escudé, J. 1996, *ApJ*, 457, L51
- Hu, E. M., Kim, T.-S., Cowie, L. L., Songaila, A., Rauch, M. 1995, *AJ*, 110, 1526
- Hui, L., Gnedin, N. Y. 1997, *MNRAS*, 292, 27
- Hui, L., Gnedin, N. Y., Zhang, Y. 1997, *ApJ*, 486, 599
- Hui, L., Rutledge, R. E. 1999, *ApJ*, 517, 541
- Kendall, M. G. & Stuart, A., 1961. *The Advanced Theory of Statistics*, Vol. 2, Griffin & Griffin, London
- Khare, P., Srianand, R., York, D. G., Green, R., Welty, D. 1997, *MNRAS*, 285, 167
- Kirkman, D., Tytler, D. 1997, *ApJ*, 484, 672
- Lu, L., Sargent, W. L. W., Womble, D. S., & Takada-Hidai, M. 1996, *ApJ*, 472, 509
- Madau, P., & Efstathiou, G. 1999, *ApJ*, 517, L9
- Miralda-Escudé, J., Cen, R., Ostriker, J. P., & Rauch, M. 1996, *ApJ*, 471, 582
- Miralda-Escudé, J., & Rees, M. J. 1994, *MNRAS*, 266, 343
- Nath, B. B., Sethi, S. K., & Shchekinov, Y. 1999, *MNRAS*, 303, 1
- Outram, P. J., Chaffee, F. H., & Carswell, R. F. 1999, *MNRAS*, submitted (astro-ph/9907042)
- Penton, S. V., Stocke, J. T., & Shull, J. M. 1999, *ApJ*, submitted
- Reimers D., Köhler S., Wisotzki L., Groote D., Rodriguez-Pascal P., & Wamsteker W., 1997, *A&A*, 327, 890

- Schaye, J., Theuns, T., Leonard, A., & Efstathiou, G. 1999, MNRAS, in press (astro-ph/9906271)
- Shull, J. M., Penton, S. V., & Stocke, J. T. 1999, Publ. Astr. Soc. Austr., Vol 16, 95
- Shull, J. M., Roberts, D., Giroux, M. L., Penton, S. V., & Fardal, M. A. 1999, AJ, 118, 1450
- Shull, J. M., Stocke, J. T., & Penton, S. 1996, AJ, 111, 72
- Theuns, T., Leonard, A., Efstathiou, G., Pearce, F. R., & Thomas, P. A. 1999, MNRAS, 301, 478
- Valageas, P., & Silk, J. 1999, A&A, 347, 1
- Weinberg, D. H., Katz, N., & Hernquist, L. 1998, ASP Conference Series, Vol. 148, eds. C. E. Woodward, J. M. Shull, H. A. Thronson, 21
- Weymann, R. J., Stern, D., Bunker, A., Spinrad, H., Chaffee, F. H., Thompson, R. I., & Storrie-Lombardi, L. J. 1999, ApJ, 505, L95
- Zhang, Y., Anninos, P., Norman, M. L., & Meiksin, A. 1997, ApJ, 485, 496